

## Clinical Policy: Phototherapy for Neonatal Hyperbilirubinemia

Reference Number: WNC.CP.131

Last Review Date: 11/2023

[Coding Implications](#)

[Revision Log](#)

See [Important Reminder](#) at the end of this policy for important regulatory and legal information.

**Note:** When state Medicaid coverage provisions conflict with the coverage provisions in this clinical policy, state Medicaid coverage provisions take precedence. Please refer to the state Medicaid manual for any coverage provisions pertaining to this clinical policy.

---

### Description

This policy details medical necessity criteria for home phototherapy for the treatment of neonatal hyperbilirubinemia. Almost all newborns will develop total serum bilirubin (TSB) levels greater than the upper limit of normal for adults, 1 mg/dL. Increasing TSB can cause jaundice, and newborns with severe hyperbilirubinemia are at risk for developing acute neurotoxicity as bilirubin crosses the blood-brain barrier. Acute bilirubin-induced neurologic dysfunction (BIND) can have chronic and permanent neurologic effects, termed kernicterus. Thus, screening for hyperbilirubinemia should be conducted on all infants prior to discharge.<sup>3</sup>

### Policy/Criteria

- I. It is the policy of WellCare of North Carolina® that conventional phototherapy in the home, applied by a single light source in the blue-green spectrum, (460-490nm), for the treatment of physiologic hyperbilirubinemia in term ( $\geq 38$  weeks gestation) infants is **medically necessary** when meeting all of the following guidelines:
  - A. Term infant status is one of the following:
    1. Previously discharged home and readmission is being considered only for hyperbilirubinemia;
    2. Infant is currently inpatient and ready for discharge except for needing treatment for elevated bilirubin;
  - B. The infant is feeding well, is active, and clinically well;
  - C. If breastfeeding or chest feeding, lactation support from a qualified professional has been offered;
  - D. A primary provider is willing to manage home care with established follow-up within the next 12-24 hours;
  - E.  $\geq 48$  hours old;
  - F. An LED-based phototherapy device will be available in the home without delay;
  - G. No previous phototherapy;
  - H. TSB will be measured daily;
  - I. Infant has none of the following risk factors:
    1. Isoimmune hemolytic disease (i.e., positive direct antiglobulin test), glucose-6-phosphate dehydrogenase (G6PD), or other hemolytic disease;

**CLINICAL POLICY WNC.CP.131  
PHOTOTHERAPY FOR NEONATAL HYPERBILIRUBINEMIA**

2. Hypoxic Ischemia Encephalopathy (HIE)/Asphyxia;
3. Temperature instability;
4. Sepsis;
5. Acidosis;
6. Albumin < 3.0 g/dL (if measured);
7. Birth weight < 2500g;
8. Significant cephalohematoma or bruising;
9. Weight loss >10%;
10. Elevated direct reacting/conjugated bilirubin;
11. Jaundice appearance in first 24 hours of life;
12. Laboratory or clinical evidence of hypothyroidism;
13. Significant clinical instability in the previous 24 hours;
14. Clinical history of a parent or sibling requiring phototherapy or exchange transfusion;
15. Exclusive breastfeeding or chest feeding with suboptimal intake ( $\geq 10\%$  weight loss);
16. Down syndrome;
17. Macrosomic infant of a diabetic mother.

J. TSB is within the levels noted in Table 1 below<sup>1</sup> :

Table 1. Acceptable TSB levels for home phototherapy in infants without risk factors, by age

Age	TSB Level
24-36 hours	$\leq 11$ mg/dL
36-48 hours	$\leq 14$ mg/dL
48-60 hours	$\leq 15$ mg/dL
60-72 hours	$\leq 16$ mg/dL
>72 hours	$\leq 17$ mg/dL

**\*Note:** The TSB home phototherapy table above allows for conservative TSB levels to align with the lower age limit in hours provided in the age ranges for inpatient criteria for hyperbilirubinemia (see section II).

**II.** It is the policy of WellCare of North Carolina® that when criteria for home phototherapy are met and home phototherapy is available in a timely fashion, inpatient phototherapy for hyperbilirubinemia is **not medically necessary** unless documentation of extenuating circumstances (including, but not limited to, expected lack of compliance with therapy at home) is provided.

**\* Note:**

- Infants should be admitted for inpatient phototherapy if the TSB concentration is more than 1 mg/dL above the AAP guidelines phototherapy treatment threshold in the hyperbilirubinemia risk calculator at

## CLINICAL POLICY WNC.CP.131

### PHOTOTHERAPY FOR NEONATAL HYPERBILIRUBINEMIA

<https://peditools.org/bili2022/index.php>. The values in Table 1 above offer phototherapy at levels consistent with the AAP statement that phototherapy can be offered below the AAP treatment threshold per the provider's discretion.

- Further criteria for inpatient phototherapy for hyperbilirubinemia can be found in clinical decision support tools.

**III.** It is the policy of WellCare of North Carolina<sup>®</sup> that other treatment for hyperbilirubinemia, including inpatient phototherapy (when not meeting criteria for home phototherapy per this policy) and exchange transfusion, is **medically necessary** when meeting the most current version of the relevant nationally recognized decision support tools.

#### Background

Efforts to reduce kernicterus include prevention and management of hyperbilirubinemia. Preventive strategies focus on identifying at-risk infants and beginning preventive therapeutic interventions as needed, usually through universal screening of all neonates for hyperbilirubinemia, which may be performed by measurement of total serum bilirubin (TSB) or by use of a transcutaneous device to obtain a Transcutaneous bilirubin (TcB) level.<sup>2</sup>

G6PD deficiency is now recognized as one of the most significant causes of hyperbilirubinemia leading to kernicterus. Identifying neonates with G6PD deficiency is challenging, so knowledge of certain risk factors for this deficiency can lead to improved health outcomes. G6PD deficiency is more common in males because it is a sex-linked recessive gene located on the X chromosome, and males only have one X chromosome. G6PD deficiency is prevalent in populations with genetic ancestry from Sub-Saharan Africa, Middle East, Mediterranean, Arabian Peninsula, and Southeast Asia. Additionally, 13% of African American males and 4% of African American females have G6PD deficiency.<sup>1</sup>

Phototherapy is considered first-line treatment for neonatal hyperbilirubinemia, defined as TSB > 95th percentile on the hour-specific Bhutani nomogram for infants  $\geq 35$  weeks gestational age (GA).<sup>1</sup> Phototherapy has been used widely for over 60 years and has been associated with few adverse events in term infants. Phototherapy decreases or reduces the rate of rise of bilirubinemia in almost all cases, regardless of the cause.<sup>2</sup> It also reduces the risk that TSB will reach the level associated with increased risk of kernicterus and that at which exchange transfusion is recommended.

Some infants are more likely to be readmitted for treatment of hyperbilirubinemia after discharge from the birth hospitalization. Infants discharged in the first two days after birth were more likely to be readmitted for jaundice compared with infants who stayed longer than three days, an association that decreased with increasing GA.<sup>7</sup> Other risk factors for hyperbilirubinemia include vaginal delivery, exclusively breastfeeding at discharge, primiparous mother, maternal age less than 20 years old, mother with an Asian country of birth, and higher TSB relative to the treatment threshold at phototherapy initiation.<sup>6,7</sup>

## CLINICAL POLICY WNC.CP.131

### PHOTOTHERAPY FOR NEONATAL HYPERBILIRUBINEMIA

Phototherapy works by using photons from light to alter bilirubin molecules in the superficial capillaries into water-soluble, non-neurotoxic molecules and reducing unconjugated TB levels.<sup>3</sup> Conventional phototherapy is delivered by a single light source. The preferred treatment is intensive phototherapy delivered by irradiance in the blue-green spectrum (wavelengths of approximately 460–490 nm) of at least 30  $\mu\text{W}/\text{cm}^2$  per nm (measured at the infant's skin directly below the center of the phototherapy unit) and delivered to as much of the infant's surface area as possible.<sup>3,4</sup> Conventional phototherapy may be delivered in the hospital setting or in the home setting.<sup>5</sup>

Home phototherapy can be less disruptive to the family and is appropriate for otherwise healthy, term infants without hemolysis and other risk factors, who have TB levels 2 to 3 mg/dL below the recommended threshold level for initiation of hospital phototherapy, are feeding well, and can be closely followed.<sup>3</sup> Per the updated 2022 clinical practice guidelines from the American Academy of Pediatrics (AAP), home phototherapy is an option that can be started at a lower threshold, such as 2 mg/dL below the phototherapy threshold, to reduce the risk of hospital readmission.<sup>1</sup>

During phototherapy, infants should be placed on their backs and fully exposed to the light with the exception of a diaper. Their eyes should be shielded with an opaque blindfold with attention given to prevent the blindfold from covering the nose or sliding off the eyes. 3

#### *American Academy of Pediatrics (AAP) <sup>1</sup>*

In 2022 the AAP published updated clinical practice guidelines that are meant to replace the 2004 clinical guidelines concerning the assessment and treatment of neonatal hyperbilirubinemia in infants  $\geq 35$  weeks.<sup>1</sup> The 2022 AAP guidelines focus on recommendations for when infants should have a direct antiglobulin test (DAT) and blood type testing; implementation of care practices that promote evidence-based breastfeeding support that is family-centered; recommendation against providing the infant with water or dextrose water as an oral supplementation to prevent hyperbilirubinemia or to decrease bilirubin levels; importance of assessing for glucose-6-phosphate dehydrogenase (G6PD) deficiency; assessment for hyperbilirubinemia neurotoxicity risk factors; recommendations for when total serum bilirubin (TSB) or transcutaneous bilirubin (TcB) should be measured; recommendations for phototherapy treatment. The 2022 AAP guidelines address the issues of prevention, risk assessment, monitoring, and treatment of neonatal hyperbilirubinemia in infants  $\geq 35$  weeks.<sup>1</sup>

#### *National Institute for Health and Care Excellence (NICE)<sup>87</sup>*

NICE guidelines cover diagnosing and treating jaundice in order to detect and prevent very high levels of bilirubin. They provide consensus-based thresholds for when phototherapy and exchange transfusion should be initiated by age in hours.<sup>8</sup>

#### *United States Preventive Services Task Force (USPSTF)<sup>98</sup>*

The USPSTF stated there was insufficient evidence to make recommendations regarding screening for hyperbilirubinemia for infants  $\geq 35$  weeks. They note that risk factors for hyperbilirubinemia include family history of neonatal jaundice, exclusive breastfeeding, bruising, cephalohematoma, ethnicity (Asian or Black), maternal age older than 25 years, male

**CLINICAL POLICY WNC.CP.131**  
**PHOTOTHERAPY FOR NEONATAL HYPERBILIRUBINEMIA**

sex, glucose-6-phosphate dehydrogenase deficiency, and gestational age less than 38 weeks. The specific contribution of these risk factors to chronic bilirubin encephalopathy in healthy children is not well understood. Currently, the USPSTF notes this recommendation is “inactive.”<sup>9</sup>

**Coding Implications**

This clinical policy references Current Procedural Terminology (CPT®). CPT® is a registered trademark of the American Medical Association. All CPT codes and descriptions are copyrighted 2020, American Medical Association. All rights reserved. CPT codes and CPT descriptions are from the current manuals and those included herein are not intended to be all-inclusive and are included for informational purposes only. Codes referenced in this clinical policy are for informational purposes only. Inclusion or exclusion of any codes does not guarantee coverage. Providers should reference the most up-to-date sources of professional coding guidance prior to the submission of claims for reimbursement of covered services.

<b>CPT®*</b>	<b>Description</b>
<b>Codes</b>	
No applicable codes.	

<b>HCPCS ®*</b>	<b>Description</b>
<b>Codes</b>	
E0202	Phototherapy (bilirubin) light with photometer
S9098	Home visit, phototherapy services (e.g., Bili-lite), including equipment rental, nursing services, blood draw, supplies, and other services, per diem

<b>Reviews, Revisions, and Approvals</b>	<b>Reviewed Date</b>	<b>Approval Date</b>
Original approval date	03/21	05/21
Reviewed HCPCS and ICD-10-CM codes.	08/21	11/21
Annual review. Clarified in section III. that the statement “applies when not meeting criteria for home phototherapy in this policy.” Background updated with no clinical significance. References reviewed and updated.	08/22	08/22
Annual review. Changed title from “Home phototherapy...” to “Phototherapy...” Updated criteria I.D. from 24-48 hours to 12-24 hours. Updated criteria to include the following: I.E., ≥48 hours old; I.F. An LED-based phototherapy device will be available in the home without delay; I.G. No previous phototherapy; I.H. TSB will be measured daily. Criteria I.I. #1 updated to include example of positive direct antiglobulin test for isoimmune hemolytic disease and to include glucose-6-phosphate dehydrogenase (G6PD) and other hemolytic disease. Criteria I.I. #2 updated to include hypoxic ischemia encephalopathy (HIE). Significant lethargy removed from Criteria I.I. Criteria I.I. updated to include the following: #13 Significant clinical	11/22	11/22

**CLINICAL POLICY WNC.CP.131  
PHOTOTHERAPY FOR NEONATAL HYPERBILIRUBINEMIA**



Reviews, Revisions, and Approvals	Reviewed Date	Approval Date
<p>instability in the previous 24 hours; #14 Clinical history of a parent or sibling requiring phototherapy or exchange transfusion; #15 Exclusive breastfeeding with suboptimal intake (<math>\geq 10\%</math> weight loss); #16 Down syndrome; #17 Macrosomic infant of a diabetic mother. Added note below Table 1 that explains the values are conservative TSB values based on lower age range thresholds in inpatient criteria. Added clarification to II that extenuating circumstances can include lack of expected compliance with therapy at home. Added note below policy statement II stating: that infants should be admitted for inpatient phototherapy if the TSB concentration is more than 1 mg/dL above the AAP guidelines phototherapy treatment threshold per the Bili risk tool, and that table 1 is consistent with AAP guidelines allowing treatment at lower levels per provider discretion; and that clinical decision support tools provider further criteria for inpatient phototherapy treatment. Updated background to include 2022 AAP clinical practice guidelines.. References reviewed and updated. Reviewed by internal specialist and external specialist.</p>		
<p>NCHC verbiage removed from NC Guidance Verbiage.</p>	<p>04/23</p>	<p>04/23</p>
<p>Annual Review. I.C. “If the mother is breastfeeding, she has been offered lactation support from a qualified professional;” is changed to: “If breastfeeding or chestfeeding, lactation support from a qualified professional has been offered;” I.I.15 “Exclusive breastfeeding with suboptimal intake (<math>\geq 10\%</math> weight loss);” has been changed to “Exclusive breastfeeding or chestfeeding with suboptimal intake (<math>\geq 10\%</math> weight loss);” Removed ICD-10-CMS codes.</p>	<p>11/23</p>	<p>11/23</p>

**References**

1. Kemper AR, Newman TB, Slaughter JL, et al. Clinical Practice Guideline Revision: Management of Hyperbilirubinemia in the Newborn Infant 35 or More Weeks of Gestation. *Pediatrics*. 2022;150(3):e2022058859. doi:10.1542/peds.2022-058859
2. Morris BH, Oh W, Tyson JE, et al. Aggressive vs. conservative phototherapy for infants with extremely low birth weight. *New Engl J Med*. 2008;359:1885 to 86. doi: 10.1056/NEJMoa0803024
3. Wong RJ, Bhutani VK. Unconjugated hyperbilirubinemia in the newborn: Interventions. UpToDate. [www.uptodate.com](http://www.uptodate.com). Updated November 10, 2021. Accessed September 07, 2022.
4. Kumar P, Chawla D, Deorari A. Light-emitting diode phototherapy for unconjugated hyperbilirubinemia in neonates. *Cochrane Database Syst Rev*. 2011;2011(12):CD007969. Published 2011 Dec 7. doi:10.1002/14651858.CD007969.pub2
5. Snook, J., 2017. Is home phototherapy in the term neonate with physiological jaundice a feasible practice? A systematic literature review. *J Neonatal Nurs*. 2017;23(1):28 to 39.

6. Chang PW, Waite WM. Evaluation of Home Phototherapy for Neonatal Hyperbilirubinemia. *J Pediatr*. 2020;220:80 to 85. doi:10.1016/j.jpeds.2020.01.004
7. Lain SJ, Roberts CL, Bowen JR, Nassar N. Early discharge of infants and risk of readmission for jaundice. *Pediatrics*. 2015;135(2):314 to 321. doi:10.1542/peds.2014-2388
8. Jaundice in newborn babies under 28 days: Clinical guideline CG98. National Institute for Health and Care Excellence. <https://www.nice.org.uk/guidance/cg98>. Published May 19, 2010 (updated October 26, 2016). Accessed September 02, 2022.
9. US Preventive Services Task Force. Screening of infants for hyperbilirubinemia to prevent chronic bilirubin encephalopathy: US Preventive Services Task Force recommendation statement. *Pediatrics*. 2009;124(4):1172 to 1177. doi:10.1542/peds.2009-0128
10. Maisels MJ, Bhutani VK, Bogen D, Newman TB, Stark AR, Watchko JF. Hyperbilirubinemia in the newborn infant > or =35 weeks' gestation: an update with clarifications. *Pediatrics*. 2009;124(4):1193 to 1198. doi:10.1542/peds.2009-0329
11. Maisels MJ, Watchko JF, Bhutani VK, Stevenson DK. An approach to the management of hyperbilirubinemia in the preterm infant less than 35 weeks of gestation. *J Perinatol*. 2012;32(9):660 to 664. doi:10.1038/jp.2012.71
12. Morris BH, Oh W, Tyson JE, et al. Aggressive vs. conservative phototherapy for infants with extremely low birth weight. *N Engl J Med*. 2008;359(18):1885 to 1896. doi:10.1056/NEJMoa0803024
13. Bhutani VK, Wong RJ. Unconjugated hyperbilirubinemia in the preterm infant (less than 35 weeks gestation). UpToDate. [www.uptodate.com](http://www.uptodate.com). Updated December 02, 2021. Accessed September 02, 2022.
14. Johnson L, Bhutani VK, Karp K, Sivieri EM, Shapiro SM. Clinical report from the pilot USA Kernicterus Registry (1992 to 2004). *J Perinatol*. 2009;29 Suppl 1:S25 to S45. doi:10.1038/jp.2008.211
15. American Academy of Pediatrics and American College of Obstetrics and Gynecology. *Guidelines for Perinatal Care*: 8th Edition. Elk Grove Village, IL. 2012.
16. Chu L, Qiao J, Xu C. Home-Based Phototherapy Versus Hospital-Based Phototherapy for Treatment of Neonatal Hyperbilirubinemia: A Systematic Review and Meta-Analysis. *Clin Pediatr (Phila)*. 2020;59(6):588 to 595. doi:10.1177/0009922820916894
17. Wong RJ, Bhutani VK. Unconjugated hyperbilirubinemia in term and late preterm infants: Management. UpToDate. [www.uptodate.com](http://www.uptodate.com). Updated December 02, 2021. Accessed September 02, 2022.
18. Pettersson M, Eriksson M, Albinsson E, Ohlin A. Home phototherapy for hyperbilirubinemia in term neonates-an unblinded multicentre randomized controlled trial. *Eur J Pediatr*. 2021;180(5):1603 to 1610. doi:10.1007/s00431-021-03932-4

### **North Carolina Guidance**

#### *Eligibility Requirements*

- a. An eligible beneficiary shall be enrolled in the NC Medicaid Program (Medicaid is NC Medicaid program, unless context clearly indicates otherwise);
- b. Provider(s) shall verify each Medicaid beneficiary's eligibility each time a service is rendered.

## CLINICAL POLICY WNC.CP.131

### PHOTOTHERAPY FOR NEONATAL HYPERBILIRUBINEMIA

- c. The Medicaid beneficiary may have service restrictions due to their eligibility category that would make them ineligible for this service.

#### *EPSDT Special Provision: Exception to Policy Limitations for a Medicaid Beneficiary under 21 Years of Age*

- a. 42 U.S.C. § 1396d(r) [1905(r) of the Social Security Act]  
Early and Periodic Screening, Diagnostic, and Treatment (EPSDT) is a federal Medicaid requirement that requires the state Medicaid agency to cover services, products, or procedures for Medicaid beneficiary under 21 years of age if the service is medically necessary health care to correct or ameliorate a defect, physical or mental illness, or a condition [health problem] identified through a screening examination (includes any evaluation by a physician or other licensed practitioner).

This means EPSDT covers most of the medical or remedial care a child needs to improve or maintain his or her health in the best condition possible, compensate for a health problem, prevent it from worsening, or prevent the development of additional health problems.

Medically necessary services will be provided in the most economic mode, as long as the treatment made available is similarly efficacious to the service requested by the beneficiary's physician, therapist, or other licensed practitioner; the determination process does not delay the delivery of the needed service; and the determination does not limit the beneficiary's right to a free choice of providers.

EPSDT does not require the state Medicaid agency to provide any service, product or procedure:

1. that is unsafe, ineffective, or experimental or investigational.
2. that is not medical in nature or not generally recognized as an accepted method of medical practice or treatment.

Service limitations on scope, amount, duration, frequency, location of service, and other specific criteria described in clinical coverage policies may be exceeded or may not apply as long as the provider's documentation shows that the requested service is medically necessary "to correct or ameliorate a defect, physical or mental illness, or a condition" [health problem]; that is, provider documentation shows how the service, product, or procedure meets all EPSDT criteria, including to correct or improve or maintain the beneficiary's health in the best condition possible, compensate for a health problem, prevent it from worsening, or prevent the development of additional health problems.

#### **EPSDT and Prior Approval Requirements**

1. If the service, product, or procedure requires prior approval, the fact that the beneficiary is under 21 years of age does NOT eliminate the requirement for prior approval.
2. **IMPORTANT ADDITIONAL INFORMATION** about EPSDT and prior approval is found in the *NCTracks Provider Claims and*



**CLINICAL POLICY WNC.CP.131**  
**PHOTOTHERAPY FOR NEONATAL HYPERBILIRUBINEMIA**



*Billing Assistance Guide*, and on the EPSDT provider page. The Web addresses are specified below:

*NCTracks Provider Claims and Billing Assistance Guide:*

<https://www.nctracks.nc.gov/content/public/providers/provider-manuals.html>

*EPSDT provider page:* <https://medicaid.ncdhhs.gov/>

*Provider(s) Eligible to Bill for the Procedure, Product, or Service*

To be eligible to bill for the procedure, product, or service related to this policy, the provider(s) shall:

- a. meet Medicaid qualifications for participation;
- b. have a current and signed Department of Health and Human Services (DHHS) Provider Administrative Participation Agreement; and
- c. bill only for procedures, products, and services that are within the scope of their clinical practice, as defined by the appropriate licensing entity.

*Compliance*

Provider(s) shall comply with the following in effect at the time the service is rendered:

- a. All applicable agreements, federal, state and local laws and regulations including the Health Insurance Portability and Accountability Act (HIPAA) and record retention requirements; and
- b. All NC Medicaid's clinical (medical) coverage policies, guidelines, policies, provider manuals, implementation updates, and bulletins published by the Centers for Medicare and Medicaid Services (CMS), DHHS, DHHS division(s) or fiscal contractor(s).

*Claims-Related Information*

Provider(s) shall comply with the NC Tracks Provider Claims and Billing Assistance Guide, Medicaid bulletins, fee schedules, NC Medicaid's clinical coverage policies and any other relevant documents for specific coverage and reimbursement for Medicaid:

- a. Claim Type - as applicable to the service provided:  
Professional (CMS-1500/837P transaction)  
Institutional (UB-04/837I transaction)  
Unless directed otherwise, Institutional Claims must be billed according to the National Uniform Billing Guidelines. All claims must comply with National Coding Guidelines.
- b. International Classification of Diseases and Related Health Problems, Tenth Revisions, Clinical Modification (ICD-10-CM) and Procedural Coding System (PCS) - Provider(s) shall report the ICD-10-CM and Procedural Coding System (PCS) to the highest level of specificity that supports medical necessity. Provider(s) shall use the current ICD-10 edition and any subsequent editions in effect at the time of service. Provider(s) shall refer to the applicable edition for code description, as it is no longer documented in the policy.
- c. Code(s) - Provider(s) shall report the most specific billing code that accurately and completely describes the procedure, product or service provided. Provider(s) shall use the Current Procedural Terminology (CPT), Health Care Procedure Coding System (HCPCS), and UB-04 Data Specifications Manual (for a complete listing of valid revenue codes) and any subsequent editions in effect at the time of service. Provider(s) shall refer to the applicable edition for the code description, as it is no longer documented in the

## CLINICAL POLICY WNC.CP.131

### PHOTOTHERAPY FOR NEONATAL HYPERBILIRUBINEMIA

policy. If no such specific CPT or HCPCS code exists, then the provider(s) shall report the procedure, product or service using the appropriate unlisted procedure or service code.

#### *Unlisted Procedure or Service*

CPT: The provider(s) shall refer to and comply with the Instructions for Use of the CPT Codebook, Unlisted Procedure or Service, and Special Report as documented in the current CPT in effect at the time of service.

HCPCS: The provider(s) shall refer to and comply with the Instructions For Use of HCPCS National Level II codes, Unlisted Procedure or Service and Special Report as documented in the current HCPCS edition in effect at the time of service

- d. Modifiers - Providers shall follow applicable modifier guidelines.
- e. Billing Units - Provider(s) shall report the appropriate code(s) used which determines the billing unit(s).
- f. Co-payments -  
For Medicaid refer to Medicaid State Plan:  
<https://medicaid.ncdhhs.gov/get-involved/nc-health-choice-state-plan>
- g. Reimbursement - Provider(s) shall bill their usual and customary charges. For a schedule of rates, refer to: <https://medicaid.ncdhhs.gov/>.

#### **Important Reminder**

This clinical policy has been developed by appropriately experienced and licensed health care professionals based on a review and consideration of currently available generally accepted standards of medical practice; peer-reviewed medical literature; government agency/program approval status; evidence-based guidelines and positions of leading national health professional organizations; views of physicians practicing in relevant clinical areas affected by this clinical policy; and other available clinical information. The Health Plan makes no representations and accepts no liability with respect to the content of any external information used or relied upon in developing this clinical policy. This clinical policy is consistent with standards of medical practice current at the time that this clinical policy was approved. “Health Plan” means a health plan that has adopted this clinical policy and that is operated or administered, in whole or in part, by Centene Management Company, LLC, or any of such health plan’s affiliates, as applicable.

The purpose of this clinical policy is to provide a guide to medical necessity, which is a component of the guidelines used to assist in making coverage decisions and administering benefits. It does not constitute a contract or guarantee regarding payment or results. Coverage decisions and the administration of benefits are subject to all terms, conditions, exclusions and limitations of the coverage documents (e.g., evidence of coverage, certificate of coverage, policy, contract of insurance, etc.), as well as to state and federal requirements and applicable Health Plan-level administrative policies and procedures.

This clinical policy is effective as of the date determined by the Health Plan. The date of posting may not be the effective date of this clinical policy. This clinical policy may be subject to applicable legal and regulatory requirements relating to provider notification. If there is a

**CLINICAL POLICY WNC.CP.131**  
**PHOTOTHERAPY FOR NEONATAL HYPERBILIRUBINEMIA**

discrepancy between the effective date of this clinical policy and any applicable legal or regulatory requirement, the requirements of law and regulation shall govern. The Health Plan retains the right to change, amend or withdraw this clinical policy, and additional clinical policies may be developed and adopted as needed, at any time.

This clinical policy does not constitute medical advice, medical treatment or medical care. It is not intended to dictate to providers how to practice medicine. Providers are expected to exercise professional medical judgment in providing the most appropriate care, and are solely responsible for the medical advice and treatment of members/enrollees. This clinical policy is not intended to recommend treatment for members/enrollees. Members/enrollees should consult with their treating physician in connection with diagnosis and treatment decisions.

Providers referred to in this clinical policy are independent contractors who exercise independent judgment and over whom the Health Plan has no control or right of control. Providers are not agents or employees of the Health Plan.

This clinical policy is the property of the Health Plan. Unauthorized copying, use, and distribution of this clinical policy or any information contained herein are strictly prohibited. Providers, members/enrollees and their representatives are bound to the terms and conditions expressed herein through the terms of their contracts. Where no such contract exists, providers, members/enrollees and their representatives agree to be bound by such terms and conditions by providing services to members/enrollees and/or submitting claims for payment for such services.

©2018 Centene Corporation. All rights reserved. All materials are exclusively owned by Centene Corporation and are protected by United States copyright law and international copyright law. No part of this publication may be reproduced, copied, modified, distributed, displayed, stored in a retrieval system, transmitted in any form or by any means, or otherwise published without the prior written permission of Centene Corporation. You may not alter or remove any trademark, copyright or other notice contained herein. Centene® and Centene Corporation® are registered trademarks exclusively owned by Centene Corporation.